

The Parkinson's disease death rate: carbidopa and vitamin B6

Marty Hinz¹

Alvin Stein²

Ted Cole³

¹Clinical Research, NeuroResearch Clinics, Inc., Cape Coral, FL, USA;

²Stein Orthopedic Associates, Plantation, FL, USA; ³Cole Center for Healing, Cincinnati, OH, USA

Abstract: The only indication for carbidopa and benserazide is the management of L-3,4-dihydroxyphenylalanine (L-dopa)-induced nausea. Both drugs irreversibly bind to and permanently deactivate pyridoxal 5'-phosphate (PLP), the active form of vitamin B6, and PLP-dependent enzymes. PLP is required for the function of over 300 enzymes and proteins. Virtually every major system in the body is impacted directly or indirectly by PLP. The administration of carbidopa and benserazide potentially induces a nutritional catastrophe. During the first 15 years of prescribing L-dopa, a decreasing Parkinson's disease death rate was observed. Then, in 1976, 1 year after US Food and Drug Administration approved the original L-dopa/carbidopa combination drug, the Parkinson's disease death rate started increasing. This trend has continued to the present, for 38 years and counting. The previous literature documents this increasing death rate, but no hypothesis has been offered concerning this trend. Carbidopa is postulated to contribute to the increasing Parkinson's disease death rate and to the classification of Parkinson's as a progressive neurodegenerative disease. It may contribute to L-dopa tachyphylaxis.

Keywords: L-dopa, levodopa, vitamin B6, pyridoxal 5'-phosphate

Introduction

Parkinson's disease is classified as a progressive neurodegenerative disease.¹ L-3,4-dihydroxyphenylalanine (L-dopa) is the most effective treatment for Parkinson's disease.² Many patients who take it experience profound nausea, which may prevent them from reaching higher dosing values required for symptom relief.³ On May 2, 1975, the US Food and Drug Administration (FDA) approved carbidopa (MK-486),^{4,5} a drug whose only indication was the management of L-dopa-induced nausea. The Centers for Disease Control and Prevention (CDC) noted an increasing Parkinson's disease death rate. In 2003, Parkinson's disease was added to the top 15 causes of death; it entered the list as the 14th leading cause of death.⁶

The following questions are examined in this review:

1. Is carbidopa linked to the increasing Parkinson's disease death rate?
2. Are the attributes of the nutritional collapse associated with Parkinson's disease, L-dopa, and/or carbidopa being misdiagnosed as progressive neurodegeneration?
3. Is carbidopa involved in L-dopa tachyphylaxis?

Insight into these questions required the review of Parkinson's disease, relative nutritional deficiencies, pyridoxal 5'-phosphate (PLP) (the active form of vitamin B6), L-dopa, carbidopa, carbidopa's side effects, the CDC-reported Parkinson's disease death rates, the biochemistry of L-dopa-induced nausea, and of the documented alternatives to carbidopa and benserazide.

Correspondence: Marty Hinz
NeuroResearch Clinics, Inc.,
1008 Dolphin Dr., Cape Coral,
FL, USA 33904
Tel +1 218 626 2220
Fax +1 218 626 1638
Email marty@hinzmd.com

Benserazide

Benserazide and carbidopa have identical mechanisms of action and indications. Any reference to benserazide means, “Benserazide and/or its metabolite trihydroxybenzylhydrazine.”⁷ While the focus of this paper is on carbidopa, the attributes shared by benserazide are noted.

Drug nutrient perspective

The following definition for a nutrient is utilized: A nutrient is any substance that facilitates normal system function. A drug is any substance that induces abnormal system function. A nutrient may become a drug. A drug may not become a nutrient.

5-hydroxytryptophan (5-HTP) is a nutrient. When it is administered as a single agent, dopamine depletion may occur.⁸⁻²¹ If it induces dopamine depletion, then 5-HTP no longer functions as a nutrient; it is a drug. L-dopa may be administered as a nutrient. When it is administered as a single agent, serotonin depletion may occur.^{10-18,22-29} If it induces serotonin depletion, then L-dopa no longer functions as a nutrient; it is a drug.

Vitamin B6

Over 300 enzymes and proteins require PLP to function properly.³⁰ The five PLP-dependent enzymes – glutamate decarboxylase, arginine decarboxylase, histamine decarboxylase, aromatic L-amino acid decarboxylase (AADC), and sulfoalanine decarboxylase – are:

[...] unrivaled in the variety of reactions they catalyze and the highly diverse metabolic pathways they are involved in, including the conversion of amino acids, one-carbon units, biogenic amines, tetrapyrrolic compounds, and amino sugars [...] sulfur assimilation, incorporation in cysteine, biotin, and S-adenosyl methionine.³¹

L-dopa

The primary pathology demonstrated in Parkinson’s disease is progressive degeneration of the substantia nigra of the brain.¹² The primary etiology of Parkinson’s disease is postsynaptic dopamine neuron damage caused by neurotoxins. Progressive neuron damage induces the collapse of the electrical conduction that regulates fine motor control.¹² L-dopa crosses the blood–brain barrier and it is then freely synthesized to dopamine without feedback regulation. The administration of L-dopa increases synaptic dopamine levels.³²⁻⁵⁵ It is analogous to turning up the voltage; more electricity flows through the remaining viable postsynaptic neurons. Restoration of postsynaptic electrical flow optimizes regulation of fine motor control.^{10,56}

The only major advancements in Parkinson’s treatment occurred in the 1950s and involved the amino acid, L-dopa. This research was awarded the Nobel Prize in Medicine in 2000^{57,58} and the Nobel Prize in Chemistry in 2001.⁵⁹ Sweet and McDowell⁶⁰ attributed the decreasing Parkinson’s death rate that occurred between 1958 and 1975 to L-dopa (from 2.9/100,000 to 1.6/100,000 of the standard population, age-adjusted).⁶¹

The National Parkinson Foundation notes that 89% of the 1 million Parkinson’s patients in the US take L-dopa/carbidopa daily.⁶² While L-dopa is the most effective treatment, it is not usually the drug of choice.² Side effects have positioned it to be one of the last drugs started in many cases.²

Parkinson’s disease is associated with the depletion of serotonin, dopamine, norepinephrine, epinephrine, thiols (homocysteine, L-methionine, S-adenosyl-L-methionine, S-adenosyl-homocysteine, cystathione, L-cysteine, and glutathione), L-tyrosine, and L-tryptophan, and these depletions represent relative nutritional deficiencies (RNDs) where systemic nutritional synthesis requirements cannot be achieved on a normal or optimal diet.^{12,13,28,63-68}

L-dopa may induce its own unique RND and exacerbate the Parkinson’s disease RND. When administered singularly or with an improper nutrient balance, it has the ability to induce RNDs of serotonin, thiols, L-tyrosine, and L-tryptophan.^{12,13} We hypothesize that if the Parkinson’s disease patient is being evaluated for the signs and symptoms of progressive neurodegenerative disease without considering the potential for, as well as the existence and ramifications of RNDs associated with the disease itself, L-dopa, and/or carbidopa, then nutritional collapse components will erroneously be attributed to progressive neurodegeneration.

Carbidopa

Efficacy and safety concerns must be addressed before US FDA drug approval. Conceptualize a drug that has no treatment efficacy claims under US FDA guidelines. Its sole indication is the management of the side effect of nausea induced by improperly balanced nutrient administration.³ It irreversibly binds to and permanently deactivates free PLP and PLP-dependent enzymes while inducing PLP reserve pool depletion.⁷ It negatively impacts the function of over 300 enzymes and proteins.³⁰ Administering vitamin B6 counteracts its mechanism of action.³ Drugs with these attributes are being prescribed. (2S)-3-(3,4-dihydroxyphenyl)-2-hydrazino-2-methylpropanoic acid (carbidopa) is being prescribed in the US and (RS)-2-amino-3-hydroxy-N’-(2,3,

4-trihydroxybenzyl) propanehydrazide (benserazide) is being prescribed outside the US.^{2,69}

L-dopa-induced nausea is a peripheral phenomenon. Carbidopa and benserazide inhibit peripheral L-dopa metabolism by AADC.³ This increases the amount of L-dopa available to cross the blood–brain barrier. Decreasing peripheral L-dopa levels through decreased ingestion, while maintaining its levels in the central nervous system, effectively controls nausea.¹² Carbidopa and benserazide control nausea by identical mechanisms.^{2,69} Carbidopa and the active metabolite of benserazide, trihydroxybenzylhydrazine, irreversibly bind to and permanently deactivate PLP and PLP-dependent enzymes.^{18,19,70–74} Normally, a Schiff base aldamine reaction catalyzes the irreversible hydrazine binding of PLP with the core protein of AADC to produce the active enzyme.⁵ Benserazide is completely metabolized to trihydroxybenzylhydrazine before it reaches the arterial blood. Carbidopa and trihydroxybenzylhydrazine are substrate analogues endowed with irreversible substituted hydrazine function.⁷

PLP is noncovalently (reversibly) bound to approximately 300 enzymes and proteins forming the PLP reserve pool.³⁰ The molecular weight of PLP is 247.142 g/mol⁷⁵ and that of carbidopa is 244.244.⁷⁶ Carbidopa irreversibly binds to PLP in a 1:1 ratio.⁷ The recommended dietary allowance of vitamin B6 is about 1 to 2 mg/day depending on age.⁷⁷ If the lowest daily dose of carbidopa (10 mg) is administered, then the system is placed into a PLP-induced RND state on the first day.

Systemic vitamin B6 concentrations inversely correlate with mortality induced by coronary artery disease, colorectal cancer, stroke, heart failure, and atherosclerosis.^{78–83} We hypothesize that if carbidopa and benserazide significantly deplete PLP, then an increased death rate will be observed. During the first 15 years of prescribing L-dopa (1960–1975) it was administered without carbidopa, a practice that was associated with a decreasing death rate.⁶¹ On May 9, 1975, the US FDA approved carbidopa for concomitant administration with L-dopa.³ Between 1976 and 2011, there has been an increase in the general Parkinson's disease death rate. While numerous etiologies have been postulated to explain the increasing Parkinson's death rate, none have impacted this 38-year trend. Parkinson's disease is most prevalent in white males. Figures 1 and 2, when viewed together, demonstrate that the Parkinson's death rate increase is occurring across all ages, sexes, and ethnic groups.

Concomitant L-dopa/carbidopa preparations have been positioned by the manufacturers to permeate treatment to the




Figure 1 Parkinson's disease death rates in the United States between 1976 and 2011, all ages, age-adjusted, male and female.

Notes: Graph generated from multiple data sources.^{62,84,89}

point that prescription L-dopa as a single ingredient is no longer available (Table 1). Prior to 1999, three pharmaceutical companies distributed a US FDA-approved brand name prescription forms of L-dopa as a single-ingredient drug: Bendopa[®] (Valent Pharm Intl); Dopar[®] (Shire plc, St Helier, Jersey); and Larodopa[®] (Hoffman-La Roche Ltd, Basel, Switzerland).²² There is no public documentation explaining the reasoning




Figure 2 Parkinson's disease death rates in the United States between 1981–2010, in comparison to white males of all ages, and white males aged 65 years and older.

Note: Graph generated from the following data source: Centers for Disease Control and Prevention. National Center for Health Statistics. Health Data Interactive. www.cdc.gov/nchs/hdi.htm.⁸⁴

Table 1 The timeline of significant events**L-dopa/carbidopa timeline**

1958–1975: The Parkinson's disease death rate decreased from 2.9/100,000 to 1.6/100,000 and was attributed to L-dopa.⁶¹
 1967: The first four studies on the administration of a general decarboxylase inhibitor for the management of L-dopa-induced nausea were documented.⁸⁶
 1975: The original brand of L-dopa with carbidopa (Sinemet®) was approved by the US FDA.^{62,86}
 1976–2011: The Parkinson's disease death rate increased by 328.7%.^{61,84}
 1977: The first paper demonstrating significant peripheral and central PLP depletion by carbidopa was submitted for publication.⁵
 1999: Pharmaceutical companies discontinued distributing the prescription form of L-dopa (a single-ingredient drug leaving L-dopa/carbidopa combinations the only prescription options).⁸⁷
 2003: The CDC added Parkinson's disease to the top 15 causes of death; it entered at number 14.⁶
 2012: Paper that asserts that carbidopa irreversibly binds to PLP and PLP-dependent enzyme molecules was published. Prior to this, carbidopa depletion of PLP was viewed as a side event, not the mechanism of action.⁷

Abbreviations: L-dopa, L-3,4-dihydroxyphenylalanine; US FDA, United States Food and Drug Administration; PLP, pyridoxal 5'-phosphate; CDC, Centers for Disease Control and Prevention.

behind the virtually simultaneous discontinuation of these drugs by the three companies.

When these drugs were approved, each was described as a decarboxylase inhibitor. Documentation submitted in 1997 noted significant central and peripheral PLP depletion after limited ingestion time of carbidopa.⁵ Now the full mechanism of action of PLP is known, giving rise to more serious concerns. It is documented that PLP freely crosses the blood–brain barrier.⁸⁴ Carbidopa prescribing information states that it “[...] does not affect the metabolism of levodopa within the central nervous system.”⁷³ This is not correct. If PLP freely crosses the blood–brain barrier allowing a peripheral and central equilibrium to exist, then both peripheral and central PLP depletion will be induced by carbidopa and benserazide. Theoretically, complete PLP depletion of the central and peripheral systems may occur. If carbidopa-induced PLP depletion is great enough for compromise of central AADC function to occur, then there will be an impairment of central dopamine synthesis. This is a previously undocumented potential etiology of L-dopa tachyphylaxis.

Carbidopa prescribing information lacks full disclosure.³ There is no reference to PLP, PLP depletion, irreversible binding to and permanent deactivation of PLP and PLP-dependent molecules, depletion of PLP reserve pools, risks induced by PLP depletion, potential functional compromise of over 300 enzymes and proteins, or RND induction. Simply describing carbidopa and benserazide as decarboxylase

inhibitors is analogous to describing a nuclear blast as a window breaker. Chronic administration affects virtually every system in the body.³⁰

The mechanism of action of carbidopa and benserazide is PLP depletion.^{18,19,70–74} Carbidopa prescribing information only notes, “Pyridoxine hydrochloride (vitamin B6), in oral doses of 10 mg to 25 mg, may reverse the effects of levodopa by increasing the rate of aromatic amino acid decarboxylation. Carbidopa inhibits this action of pyridoxine.”⁷³ PLP depletion is an RND event. PLP can reverse the nausea control of carbidopa and ameliorate the clinical effects of L-dopa, requiring caution during concomitant administration. We hypothesize that if these drugs are stopped and then ample vitamin B6 is administered, PLP function, PLP-dependent enzyme function, and PLP reserve pools will return to normal.

The exact size of the PLP reserve pool, which is reversibly bound to about 300 enzymes and proteins, is a matter of speculation. We postulate that when normal PLP pool reserve function exists at the start of treatment, it may take 5 or more years of chronic carbidopa or benserazide ingestion, depending on the daily dosing value, before progressive clinical deterioration is demonstrated. We further postulate that without a PLP reserve pool, carbidopa or benserazide ingestion would induce PLP collapse in days.

Carbidopa side effect profile

L-dopa active ingredient products may be administered as a nutritional supplement or a drug. Nutritional supplements are generally recognized as safe (GRAS), allowing over-the-counter sales.¹² Due to side effects, carbidopa is not GRAS.³ It may induce life-threatening events including myocardial infarction, neuroleptic malignant syndrome, agranulocytosis, hemolytic and nonhemolytic anemia, gastrointestinal bleeding, thrombocytopenia, and hypokalemia (Table 2).^{12,13} While carbidopa side effects require its discontinuation when the continuation of L-dopa is indicated, nutrient products are the only option. Most physicians are unaware of the availability of this L-dopa form.

The carbidopa side effects and adverse reactions listed in Table 2 are a direct result of irreversible drug-induced PLP depletion, irreversible PLP-dependent enzyme binding, PLP reserve pool collapse, along with RND-induced collapse of serotonin and catecholamine synthesis.^{12,13,18,19,70–74,78–83} If, when equilibrated, central PLP depletion occurs as a result of peripheral PLP depletion, then central compromise, side effects, and adverse reactions are inevitable – a phenomenon not previously documented.

Table 2 Side effects and adverse reactions associated with carbidopa

Glossitis	Upper respiratory infection	Phlebitis
Leg pain		Agranulocytosis
Ataxia	Bruxism	Hemolytic and nonhemolytic anemia
Falling	Hiccups	
Gait abnormalities	Common cold	
Blepharospasm (which may be taken as an early sign of excess dosage)	Diarrhea	Rash
	Urinary tract infections	Gastrointestinal bleeding
Trismus	Urinary frequency	Duodenal ulcer
Increased tremor	Flatulence	Henoch–Schonlein purpura
Numbness	Priapism	Decreased hemoglobin and hematocrit
Muscle twitching	Pharyngeal pain	Thrombocytopenia
Peripheral neuropathy	Abdominal pain	Leukopenia
Myocardial infarction	Bizarre breathing patterns	Angioedema
Flushing	Burning sensation of tongue	Urticaria
Oculogyric crises		Pruritus
Diplopia	Back pain	Alopecia
Blurred vision	Shoulder pain	Dark sweat
Dilated pupils	Chest pain (noncardiac)	Abnormalities in alkaline phosphatase
Urinary retention	Muscle cramps	Abnormalities in SGOT (AST)
Urinary incontinence	Paresthesia	SGPT (ALT)
Dark urine	Increased sweating	Abnormal Coombs' test
Hoarseness	Syncope	Abnormal uric acid
Malaise	Orthostatic hypotension	Hypokalemia
Hot flashes	Asthenia (weakness)	Abnormalities in blood urea nitrogen
Sense of stimulation	Dysphagia	Increased creatinine
Dyspepsia	Horner's syndrome, mydriasis	increased serum LDH
Constipation	Dry mouth	Glycosuria
Palpitation	Sialorrhoea	
Fatigue	Neuroleptic malignant syndrome	

Note: Data from Hinz et al.^{12,13}

Abbreviations: SGOT, serum glutamic oxaloacetic transaminase; AST, aspartate aminotransferase; SGPT, serum glutamic pyruvic transaminase; ALT, alanine transaminase; LDH, lactate dehydrogenase.

Nutritional management of nausea

It was previously documented that L-dopa-induced nausea, along with Parkinson's disease, L-dopa-associated, and carbidopa-associated RND, is definitively controlled with properly balanced administration of the nutrient 5-HTP, along with L-tyrosine, a thiol (L-cysteine, glutathione, S-adenosylmethionine, or L-methionine), and cofactors (vitamin C, vitamin B6, and calcium carbonate), as facilitated by organic cation transporter type-2 functional status analysis.^{12,13}

AADC inhibition may be reversible or irreversible. The irreversible inhibition of AADC is the mechanism

of action whereby carbidopa and benserazide control L-dopa-induced nausea.^{3,18,19,70–74} Reversible inhibition of AADC in the competitive inhibition state is the mechanism of action whereby 5-HTP controls L-dopa-induced nausea. If 5-HTP effectively controls L-dopa-induced nausea, then carbidopa or benserazide is no longer indicated, and all detrimental effects discussed herein no longer apply. If 5-HTP is not administered in proper balance with amino acid precursors of other systems, then it will become a drug due to its depletion of dopamine.^{12,13}

Due to the increased frequency of the onset of new drug-induced side effects, carbidopa, monoamine oxidase inhibitors, and catechol-O-methyl transferase inhibitors need to be stopped as 5-HTP and other nutrients are started under the nutrient protocol. If carbidopa is administered with expectations of controlling L-dopa-induced nausea, then vitamin B6 cannot be replenished while taking the drug since PLP reverses the drug effects. If there is a patient history of carbidopa or benserazide ingestion, then vitamin B6 (100–300 mg/day) is indicated at the initiation of the nutrient protocol.

Discussion

Responsible physicians create an environment where optimal symptom control nurtures healing. Two medications with no efficacy claims have been prescribed for the iatrogenic mismanagement of a nutrient, L-dopa, turning it into a drug which depletes other systems.^{3,12,13,69} Their only indication is to alleviate nausea, a benign condition, while having the ability to profoundly compromise hundreds of system functions.^{3,69} In our opinion, use of these medications is a violation of the physician's oath to first, do no harm. These drugs can create fatal events, clinical deterioration, drug-induced sequelae, and risks where none previously existed due to profound multisystem nutritional collapse.^{12,13,18,19,70–74,78–83} Nausea induced by improper administration of the nutrient L-dopa should not be addressed with drugs whose mechanism of action is system-wide vitamin B6 RND, which is especially true when a drug-free nutrient management approach is available.

In 1941, almost 20 years before the dawn of L-dopa, Baker described a subgroup involving 25% of Parkinson's disease patients who achieved "definite objective improvement" with vitamin B6 administration.⁸⁵ In 2012, the literature noted, "Multifactorial neurological pathologies such as [...] Parkinson's disease [...] have also been correlated to inadequate intracellular levels of PLP."²⁵ Administration of carbidopa and benserazide should be contraindicated in these patients.

Conclusion

Between 1960 and 1974, the only prescription form of L-dopa available was the single-ingredient form that was associated with a decreasing death rate.⁶¹ In 1975, the original combination L-dopa/carbidopa drug (Sinemet®) was approved by the US FDA. Between 1976 and 2011, the CDC documented a progressive increase of 328.7% in Parkinson's disease deaths that crossed age, sex, and ethnic boundaries.^{61,84} In addition, no effective way has been discovered to truly stop what has been described as neurodegeneration.

The mechanism of action for carbidopa and benserazide induces irreversible binding to and permanent deactivation of PLP and PLP-dependent enzyme molecules, potentially inducing a negative impact on over 300 enzymes and proteins. Without the induction of PLP deficiency, the clinical effects of carbidopa and benserazide are not observed. It is a documented fact that these drugs may induce system-wide PLP depletion, representing an RND that is reversed with vitamin B6 administration. Administration of carbidopa may play a role in the escalating Parkinson's disease death rate, the exacerbation of symptoms exclusively attributed to progressive neurodegeneration, and L-dopa tachyphylaxis. The full list of biochemical compromises can only be speculated due to the ability of carbidopa and benserazide to induce hundreds of intertwined peripheral and central PLP function collapses, many of which may not be fully understood at present. It is illogical to assert that an increased carbidopa-induced death rate will not occur under these circumstances. In an attempt to control a benign condition (nausea – caused by the improperly balanced administration of a nutrient, L-dopa), the patient has been exposed to the devastating consequences of these drugs. While a formidable number of studies may still be needed to define all of the PLP depletion ramifications, they become unnecessary in the effective management of Parkinson's disease when the nutrient protocol is implemented, since carbidopa and benserazide are no longer indicated.

The administration of properly balanced nutrients, under a documented nutritional protocol for the definitive control of L-dopa-induced nausea, should raise no more concern with the caregiver or patient than administration of a multivitamin; all are GRAS. Physicians should fully understand the mechanism of action of the drugs they prescribe rather than relying on the described indications provided by the drug company. Efficacy concerns relating to the discontinuation of carbidopa and benserazide are unfounded since they have no efficacy; they only deal with L-dopa side effects. Before 1976, in the precarbidopa era,

ample studies were published documenting the efficacy of L-dopa without carbidopa.

Three questions are raised: 1) Is progressive neurodegeneration observed with Parkinson's disease intrinsic to the disease, or may some symptoms be attributed to carbidopa or benserazide-induced RND? 2) Does iatrogenic drug-induced poisoning, which may result in irreversible binding to and permanent deactivation of PLP and PLP-dependent enzyme molecules throughout the system, play a role in the increasing death rate noted by the CDC since 1976? 3) Is carbidopa or benserazide potentially involved in L-dopa tachyphylaxis? If the answer to these questions is yes, or even maybe, a greater focus on nutrition is indicated while discontinuing drugs such as carbidopa or benserazide. The doctrine of *res ipsa loquitur* (the thing speaks for itself) applies.

There is much fertile ground presented here for furthering this research started in 1997. The authors encourage continued investigation, along with dialogue, into the ramifications of carbidopa and benserazide use and the known RNDs that plague the Parkinson's disease patient.

Disclosure

MH discloses his relationship with DBS Labs, Inc. and NeuroResearch Clinics, Inc. The other authors report no conflicts of interest in this work.

References

1. National Parkinson Foundation [webpage on the Internet]. What is Parkinson's disease? Miami, FL: National Parkinson Foundation; 2014. Available from: <http://www.parkinson.org/parkinson-s-disease/pd-101/what-is-parkinson-s-disease>. Accessed July 19, 2014.
2. Mayo Clinic [webpage on the Internet]. Diseases and conditions: Parkinson's disease. Rochester, MN: Mayo Clinic; 2014. Available from: <http://www.mayoclinic.org/diseases-conditions/parkinsons-disease/basics/treatment/con-20028488>. Accessed July 19, 2014.
3. SINEMET CR (carbidopa and levodopa) tablet, extended release [prescribing information]. Whitehouse Station, NJ: Merck & Co, Inc.; 2014. Available from: <http://dailymed.nlm.nih.gov/dailymed/lookup.cfm?setid=69e575b9-f8a5-494f-b736-2520ef505cb0>. Accessed July 1, 2014.
4. US Food and Drug Administration [webpage on the Internet]. Drugs@FDA: FDA approved drug products. Silver Spring, MD: US Food and Drug Administration; 2014. Available from: <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.DrugDetails>. Accessed July 19, 2014.
5. Airoldi L, Watkins CJ, Wiggins JF, Wurtman RJ. Effect of pyridoxine on the depletion of tissue pyridoxal phosphate by carbidopa. *Metabolism*. 1978;27(7):771–779.
6. Hoyert DL, Heron MP, Murphy SL, Kung H. *National Vital Statistics Report. Deaths: Final Data for 2003*. Hyattsville, MD: National Center for Health Statistics; 2006. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr54/nvsr54_13.pdf. Accessed July 1, 2014.
7. Daidone F, Montioli R, Paiardini A, et al. Identification by virtual screening and in vitro testing of human DOPA decarboxylase inhibitors. *PLoS One*. 2012;7(2):e31610.

8. Andrews DW, Patrick RL, Barchas JD. The effects of 5-hydroxytryptophan and 5-hydroxytryptamine on dopamine synthesis and release in rat brain striatal synaptosomes. *J Neurochem*. 1978;30(2):465–470.
9. Awazi N, Guldberg HC. On the interaction of 5-hydroxytryptophan and 5-hydroxytryptamine with dopamine metabolism in the rat striatum. *Naunyn Schmiedeberg's Arch Pharmacol*. 1978;303(1):63–72.
10. Hinz M. Depression. In: Kohlstadt I, editor. *Food and Nutrients in Disease Management*. Baton Rouge, FL: CRC Press; 2009:465–481.
11. Hinz M, Stein A, Uncini T. The dual-gate lumen model of renal monoamine transport. *Neuropsychiatr Dis Treat*. 2010;6:387–392.
12. Hinz M, Stein A, Uncini T. Amino acid management of Parkinson's disease: a case study. *Int J Gen Med*. 2011;4:165–174.
13. Hinz M, Stein A, Uncini T. Relative nutritional deficiencies associated with centrally acting monoamines. *Int J Gen Med*. 2012;5:413–430.
14. Hinz M, Stein A, Uncini T. APRESS: apical regulatory super system, serotonin, and dopamine interaction. *Neuropsychiatr Dis Treat*. 2011;7:457–463.
15. Stein A, Hinz M, Uncini T. Amino acid-responsive Crohn's disease: a case study. *Clin Exp Gastroenterol*. 2010;3:171–177.
16. Hinz M, Stein A, Neff R, Weinberg R, Uncini T. Treatment of attention deficit hyperactivity disorder with monoamine amino acid precursors and organic cation transporter assay interpretation. *Neuropsychiatr Dis Treat*. 2011;7:31–38.
17. Hinz M, Stein A, Trachte G, Uncini T. Neurotransmitter testing of the urine: a comprehensive analysis. *Open Access J Urol*. 2010;2: 177–183.
18. Hinz M, Stein A, Uncini T. A pilot study differentiating recurrent major depression from bipolar disorder cycling on the depressive pole. *Neuropsychiatr Dis Treat*. 2010;6:741–747.
19. Zhelyaskov DK, Levitt M, Udenfriend S. Tryptophan derivatives as inhibitors of tyrosine hydroxylase in vivo and in vitro. *Mol Pharmacol*. 1968;4(5):445–451.
20. Ng LK, Chase TN, Colburn RW, Kopin IJ. Release of (3 H)dopamine by L-5-hydroxytryptophan. *Brain Res*. 1972;45(2):499–502.
21. Stamford JA, Kruk ZL, Millar J. Striatal dopamine terminals release serotonin after 5-HTP pretreatment: in vivo voltammetric data. *Brain Res*. 1990;515(1–2):173–180.
22. Ritvo ER, Yuwiler A, Geller E, et al. Effects of L-dopa in autism. *J Autism Child Schizophr*. 1971;1(2):190–205.
23. Wuerthele SM, Moore KE. Studies on the mechanisms of L-dopa-induced depletion of 5-hydroxytryptamine in the mouse brain. *Life Sci*. 1977;20(10):1675–1680.
24. Borah A, Mohanakumar KP. Long-term L-DOPA treatment causes indiscriminate increase in dopamine levels at the cost of serotonin synthesis in discrete brain regions of rats. *Cell Mol Neurobiol*. 2007;27(8):985–996.
25. Karobath M, Díaz JL, Huttunen MO. The effect of L-dopa on the concentrations of tryptophan, tyrosine and serotonin in rat brain. *Eur J Pharmacol*. 1971;14(4):393–396.
26. García NH, Berndt TJ, Tyce GM, Knox FG. Chronic oral L-DOPA increases dopamine and decreases serotonin excretions. *Am J Physiol*. 1999;277(5 Pt 2):R1476–R1480.
27. Carta M, Carlsson T, Kirik D, Björklund A. Dopamine released from 5-HT terminals is the cause of L-DOPA-induced dyskinesia in parkinsonian rats. *Brain*. 2007;130(Pt 7):1819–1833.
28. Carta M, Carlsson T, Muñoz A, Kirik D, Björklund A. Serotonin–dopamine interaction in the induction and maintenance of L-DOPA-induced dyskinesias. *Prog Brain Res*. 2008;172:465–478.
29. Everett GM, Borcherding JW. L-dopa: effect on concentrations of dopamine, norepinephrine, and serotonin in brains of mice. *Science*. 1970;168(3933):849–850.
30. UniProt [webpage on the Internet]. UniProt. UniProt Consortium; 2014. Available from: <http://www.uniprot.org/uniprot/?query=pyridoxal+AND+organism%3A%22Homo+sapiens+%5B9606%5D%22&sort=score>. Accessed July 4, 2014.
31. Paiardini A, Contestabile R, Buckle AM, Cellini B. PLP-dependent enzymes. *Biomed Res Int*. 2014;2014:856076.
32. Nicotra A, Parvez S. Apoptotic molecules and MPTP-induced cell death. *Neurotoxicol Teratol*. 2002;24(5):599–605.
33. Fangman A, O'Malley WE. L-dopa and the patient with Parkinson's disease. *Am J Nurs*. 1969;69(7):1455–1457.
34. Srimal RC, Dhawan BN. An analysis of methylphenidate induced gnawing in guinea pigs. *Psychopharmacologia*. 1970;18(1):99–107.
35. Leon AS, Spiegel HE, Thomas G, Abrams WB. Pyridoxine antagonism of levodopa in parkinsonism. *JAMA*. 1971;218(13):1924–1927.
36. Smythe GA, Edwards G, Graham P, Lazarus L. Biochemical diagnosis of pheochromocytoma by simultaneous measurement of urinary excretion of epinephrine and norepinephrine. *Clin Chem*. 1992;38(4):486–492.
37. Verde G, Oppizzi G, Colussi G, et al. Effect of dopamine infusion on plasma levels of growth hormone in normal subjects and in agromegalic patients. *Clin Endocrinol (Oxf)*. 1976;5(4):419–423.
38. Weiner RI, Ganong WF. Role of brain monoamines and histamine in regulation of anterior pituitary secretion. *Physiol Rev*. 1978;58(4):905–976.
39. Mason LJ, Cojocaru TT, Cole DJ. Surgical intervention and anesthetic management of the patient with Parkinson's disease. *Int Anesthesiol Clin*. 1996;34(4):133–150.
40. Volkow ND, Fowler JS, Gatley SJ, et al. PET evaluation of the dopamine system of the human brain. *J Nucl Med*. 1996;37(7):1242–1256.
41. Checkley SA. Neuroendocrine tests of monoamine function in man: a review of basic theory and its application to the study of depressive illness. *Psychol Med*. 1980;10(1):35–53.
42. Nishino T, Lahiri S. Effects of dopamine on chemoreflexes in breathing. *J Appl Physiol Respir Environ Exerc Physiol*. 1981;50(4):892–897.
43. Pollock JD, Rowland N. Peripherally administered serotonin decreases food intake in rats. *Pharmacol Biochem Behav*. 1981;15(2): 179–183.
44. Greenamyre JT. Glutamate–dopamine interactions in the basal ganglia: relationship to Parkinson's disease. *J Neural Transm Gen Sect*. 1993; 91(2–3):255–269.
45. Ward DS, Bellville JW. Effect of intravenous dopamine on hypercapnic ventilatory response in humans. *J Appl Physiol Respir Environ Exerc Physiol*. 1983;55(5):1418–1425.
46. Morton JJ, Connell JM, Hughes MJ, Inglis GC, Wallace EC. The role of plasma osmolality, angiotensin II and dopamine in vasopressin release in man. *Clin Endocrinol (Oxf)*. 1985;23(2):129–138.
47. Seri I, Tulassay T, Kiszal J, et al. Effect of low-dose dopamine infusion on prolactin and thyrotropin secretion in preterm infants with hyaline membrane disease. *Biol Neonate*. 1985;47(6):317–322.
48. Al-Damluji S, Rees LH. Effects of catecholamines on secretion of adrenocorticotrophic hormone (ACTH) in man. *J Clin Pathol*. 1987;40(9):1098–1107.
49. Hoffman B, Lefkowitz R. Catecholamines and sympathomimetic drugs. In: Gillman AG, Rall TW, Nies AS, Taylor P, editors. *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. New York, NY: Pergamon Press; 1990.
50. Levein NG, Thörn SE, Wattwil M. Dopamine delays gastric emptying and prolongs oro-caecal transit time in volunteers. *Eur J Anaesthesiol*. 1999;16(4):246–250.
51. Bell DG, McLellan TM, Sabiston CM. Effect of ingesting caffeine and ephedrine on 10-km run performance. *Med Sci Sports Exerc*. 2002;34(2):344–349.
52. Bergerot A, Storer RJ, Goadsby PJ. Dopamine inhibits trigeminovascular transmission in the rat. *Ann Neurol*. 2007;61(3):251–262.
53. Scanlon MF, Weightman DR, Shale DJ, et al. Dopamine is a physiological regulator of thyrotrophin (TSH) secretion in normal man. *Clin Endocrinol (Oxf)*. 1979;10(1):7–15.
54. Allen GF, Land JM, Heales SJ. A new perspective on the treatment of aromatic L-amino acid decarboxylase deficiency. *Mol Genet Metab*. 2009;97(1):6–14.
55. Rubí B, Maechler P. Minireview: new roles for peripheral dopamine on metabolic control and tumor growth: let's seek the balance. *Endocrinology*. 2010;151(12):5570–5581.

56. Hinz M, Stein A, Uncini T. The discrediting of the monoamine hypothesis. *Int J Gen Med*. 2012;5:135–142.
57. Carlsson A. *A Half-Century of Neurotransmitter Research: Impact on Neurology and Psychiatry*. Nobel Lecture, December 8, 2000. Available from: http://www.nobelprize.org/nobel_prizes/medicine/laureates/2000/carlsson-lecture.pdf. Accessed July 4, 2014.
58. Kandel ER. *The Molecular Biology of Memory Storage: A Dialog Between Genes and Synapses*. Nobel Lecture, December 8, 2000. Available from: http://www.nobelprize.org/nobel_prizes/medicine/laureates/2000/kandel-lecture.pdf. Accessed July 4, 2014.
59. The Royal Swedish Academy of Sciences. *Advanced Information on the Nobel Prize in Chemistry 2001*. Stockholm, Sweden: The Royal Swedish Academy of Sciences; 2001. Available from: http://www.nobelprize.org/nobel_prizes/chemistry/laureates/2001/advanced-chemistryprize2001.pdf. Accessed July 4, 2014.
60. Sweet RD, McDowell FH. Five years' treatment of Parkinson's disease with levodopa. Therapeutic results and survival of 100 patients. *Ann Intern Med*. 1975;83(4):456–463.
61. Murphy SL, Xu JQ, Kochanek KD. Deaths: Final data for 2010. National vital statistics reports; vol 61 no 4. Hyattsville, MD: National Center for Health Statistics. 2013. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61_04.pdf. Accessed July 1, 2014.
62. National Parkinson Foundation. *The National Parkinson Foundation's Help-line Speaks: Lessons from the 2011 Sinemet Shortage*. Miami, FL: National Parkinson Foundation; 2012. Available from: http://www.parkinson.org/Files/PDFs/NPF-Content-Documents/White-Papers/NPF466_-2011-Sinemet-Shortage_WhitePaper_Full-art. Accessed July 1, 2014.
63. Mones RJ, Elizan TS, Siegel GJ. Analysis of L-dopa induced dyskinesias in 51 patients with Parkinsonism. *J Neurol Neurosurg Psychiatry*. 1971;34(6):668–673.
64. Chase TN. Serotonergic mechanisms in Parkinson's disease. *Arch Neurol*. 1972;27(4):354–356.
65. Busch AE, Karbach U, Miska D, et al. Human neurons express the polyspecific cation transporter hOCT2, which translocates monoamine neurotransmitters, amantadine, and memantine. *Mol Pharmacol*. 1998;54(2):342–352.
66. Mayeux R, Stern Y, Williams JB, Cote L, Frantz A, Dyrenfurth I. Clinical and biochemical features of depression in Parkinson's disease. *Am J Psychiatry*. 1986;143(6):756–759.
67. Chan-Palay V, Höchli M, Jentsch B, Leonard B, Zetzsche T. Raphe serotonin neurons in the human brain stem in normal controls and patients with senile dementia of the Alzheimer type and Parkinson's disease: relationship to monoamine oxidase enzyme localization. *Dementia*. 1992;3(5–6):253–269.
68. Charlton CG, Mack J. Substantia nigra degeneration and tyrosine hydroxylase depletion caused by excess S-adenosylmethionine in the rat brain. Support for an excess methylation hypothesis for parkinsonism. *Mol Neurobiol*. 1994;9(1–3):149–161.
69. Roche. Modpar [prescribing information]. Dee Why, Australia: Roche Products Pty Limited; 2010. Available from: http://www.roche-australia.com/content/dam/internet/corporate/roche/en_AU/files/central_nervous_agents/modpar-pi.pdf. Accessed July 20, 2014.
70. Bertoldi M. Mammalian Dopa decarboxylase: structure, catalytic activity and inhibition. *Arch Biochem Biophys*. 2014;546:1–7.
71. Wu F, Christen P, Gehring H. A novel approach to inhibit intracellular vitamin B6-dependent enzymes: proof of principle with human and plasmodium ornithine decarboxylase and human histidine decarboxylase. *FASEB J*. 2011;25(7):2109–2122.
72. Cellini B, Montioli R, Oppici E, Voltattorni CB. Biochemical and computational approaches to improve the clinical treatment of dopa decarboxylase-related diseases: an overview. *Open Biochem J*. 2012;6:131–138.
73. Bartlett MG. Biochemistry of the water soluble vitamins: a lecture for first year pharmacy students. *Am J Pharm Educ*. 2003; 67(2):Article 64.
74. Palfreyman MG, Danzin C, Bey P, et al. Alpha-difluoromethyl DOPA, a new enzyme-activated irreversible inhibitor of aromatic L-amino acid decarboxylase. *J Neurochem*. 1978;31(4):927–932.
75. DrugBank [webpage on the Internet]. Pyridoxal phosphate. DrugBank; 2013. Available from: <http://www.drugbank.ca/drugs/DB00114>. Accessed July 20, 2014.
76. DrugBank [webpage on the Internet]. Carbidopa. DrugBank; 2013. Available from: <http://www.drugbank.ca/drugs/DB00190>. Accessed July 2014.
77. National Institutes of Health [webpage on the Internet]. *Vitamin B6: Dietary Supplement Fact Sheet*. Bethesda, MD: National Institutes of Health; 2011. Available from: <http://ods.od.nih.gov/factsheets/VitaminB6-HealthProfessional/>. Accessed July 20, 2014.
78. Jansen MC, Bueno-de-Mesquita HB, Buzina R, et al. Dietary fiber and plant foods in relation to colorectal cancer mortality: the Seven Countries Study. *Int J Cancer*. 1999;81(2):174–179.
79. Robinson K, Arheart K, Refsum H, et al. Low circulating folate and vitamin B6 concentrations: risk factors for stroke, peripheral vascular disease, and coronary artery disease. European COMAC Group. *Circulation*. 1998;97(5):437–443.
80. Cui R, Iso H, Date C, Kikuchi S, Tamakoshi A; Japan Collaborative Cohort Study Group. Dietary folate and vitamin b6 and B12 intake in relation to mortality from cardiovascular diseases: Japan collaborative cohort study. *Stroke*. 2010;41(6):1285–1289.
81. Medrano MJ, Sierra MJ, Almazán J, Olalla MT, López-Abente G. The association of dietary folate, B6, and B12 with cardiovascular mortality in Spain: an ecological analysis. *Am J Public Health*. 2000;90(10):1636–1638.
82. Schnyder G, Roffi M, Flammer Y, Pin R, Hess OM. Effect of homocysteine-lowering therapy with folic acid, vitamin B12, and vitamin B6 on clinical outcome after percutaneous coronary intervention: the Swiss Heart study: a randomized controlled trial. *JAMA*. 2002;288(8):973–979.
83. Nygård O, Nordrehaug JE, Refsum H, Ueland PM, Farstad M, Vollset SE. Plasma homocysteine levels and mortality in patients with coronary artery disease. *N Engl J Med*. 1997;337(4):230–236.
84. Centers for Disease Control and Prevention [webpage on the Internet]. Health data interactive. Atlanta, GA: Centers for Disease Control and Prevention; 2014. Available from: <http://www.cdc.gov/nchs/hdi.htm>. Accessed July 1, 2014.
85. Baker AB. Treatment of paralysis agitans with vitamin B6 (pyridoxine hydrochloride). *JAMA*. 1941;116(22):2484–2487.
86. Giardina G, Montioli R, Gianni S, et al. Open conformation of human DOPA decarboxylase reveals the mechanism of PLP addition to Group II decarboxylases. *Proc Natl Acad Sci USA*. 2011;108(51):20514–20519.
87. Barbeau A. The pathogenesis of Parkinson's disease: a new hypothesis. *Can Med Assoc J*. 1962;87:802–807.
88. US Food and Drug Administration [webpage on the Internet]. Drugs@FDA: FDA approved drug products. Silver Spring, MD: US Food and Drug Administration; 2014. Available from: <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm>. Accessed July 2014.
89. Hoyert DL, Xu JQ. Deaths: Preliminary data for 2011. National vital statistics reports; vol 61 no 6. Hyattsville, MD: National Center for Health Statistics. 2012.

Clinical Pharmacology: Advances and Applications**Dovepress****Publish your work in this journal**

Clinical Pharmacology: Advances and Applications is an international, peer-reviewed, open access journal publishing original research, reports, reviews and commentaries on all areas of drug experience in humans. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.

Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/clinical-pharmacology-advances-and-applications-journal>